

Fortekor Flavour 5 mg tablets for cats and dogs

Species:Cats, Dogs

Therapeutic indication:Pharmaceuticals: Cardiovascular and respiratory preparations, Renal preparations

Active ingredient:Benazepril Hydrochloride

Product:Fortekor® Flavour 5mg tablets for cats and dogs

Product index:Fortekor Flavour 5mg tablets for cats and dogs

Qualitative and quantitative composition

Each tablet contains:

Active substance: Benazepril hydrochloride 5 mg

For the full list of excipients, see Pharmaceutical particulars section.

Pharmaceutical form

Tablets.

Beige to light brown, ovaloid, divisible, tablet scored on both sides.

The tablets can be divided into halves.

Clinical particulars

Target species

Dogs and cats.

Indications for use, specifying the target species

Dogs:

Treatment of congestive heart failure.

Cats:

Reduction of proteinuria associated with chronic kidney disease.

Contraindications

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

Do not use in cases of hypotension, hypovolaemia, hyponatraemia or acute renal failure.

Do not use in cases of cardiac output failure due to aortic or pulmonary stenosis.

Do not use during pregnancy or lactation (section below).

Special warnings for each target species

None.

Special precautions for use

Special precautions for use in animals

No evidence of renal toxicity of the veterinary medicinal product has been observed (in dogs or cats) during clinical trials, however, as is routine in cases of chronic kidney disease, it is recommended to monitor plasma creatinine, urea and erythrocyte counts during therapy.

The efficacy and safety of FORTEKOR has not been established in dogs and cats below 2.5 kg body weight.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use.

In case of accidental oral ingestion, seek medical advice immediately and show the label or the package leaflet to the physician.

Pregnant women should take special care to avoid accidental oral exposure because angiotensin converting enzyme (ACE) inhibitors have been found to affect the unborn child during pregnancy in humans.

Adverse reactions (frequency and seriousness)

In double-blind clinical trials in dogs with congestive heart failure, FORTEKOR Flavour was well tolerated with an incidence of adverse reactions lower than observed in placebo-treated dogs.

A small number of dogs may exhibit transient vomiting, incoordination or signs of fatigue.

In cats and dogs with chronic kidney disease, FORTEKOR Flavour may increase plasma creatinine concentrations at the start of therapy. A moderate increase in plasma creatinine concentrations following administration of ACE inhibitors is compatible with the reduction in glomerular hypertension induced by these agents, and is therefore not necessarily a reason to stop therapy in the absence of other signs.

FORTEKOR Flavour may increase food consumption and body weight in cats.

Emesis, anorexia, dehydration, lethargy and diarrhoea have been reported in rare occasions in cats.

Use during pregnancy, lactation or lay

The safety of FORTEKOR Flavour has not been established in breeding, pregnant or lactating dogs and cats.

Pregnancy

Do not use during pregnancy or lactation.

Benazepril Reduced ovary/oviduct weights in cats when administered daily at 10 mg/kg body weight for 52 weeks.

Laboratory studies in rats have shown evidence of embryotoxic effects (foetal urinary

tract malformation) at maternally non-toxic doses.

Interaction with other medicinal products and other forms of interaction

In dogs with congestive heart failure, FORTEKOR Flavour has been given in combination with digoxin, diuretics, pimobendan and anti-arrhythmic veterinary medicinal products without demonstrable adverse interactions.

In humans, the combination of ACE inhibitors and Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) can lead to reduced anti-hypertensive efficacy or impaired renal function. The combination of FORTEKOR Flavour and other anti-hypertensive agents (e.g. calcium channel blockers, β -blockers or diuretics), anaesthetics or sedatives may lead to additive hypotensive effects. Therefore, concurrent use of NSAIDs or other medications with a hypotensive effect should be considered with care. Renal function and signs of hypotension (lethargy, weakness etc) should be monitored closely and treated as necessary.

Interactions with potassium preserving diuretics like spironolactone, triamterene or amiloride cannot be ruled out. It is recommended to monitor plasma potassium levels when using FORTEKOR Flavour in combination with a potassium sparing diuretic because of the risk of hyperkalaemia.

Amounts to be administered and administration route

FORTEKOR Flavour should be given orally once daily, with or without food. The duration of treatment is unlimited.

FORTEKOR Flavour 5 mg tablets are flavoured and are taken voluntarily by most dogs and cats.

Dogs:

FORTEKOR Flavour should be administered orally at a minimum dose of 0.25 mg (range 0.25-0.5) benazepril hydrochloride/kg body weight once daily, according to the following table:

Weight of dog (kg)	Fortekor® Flavour 5 mg	
	Standard Dose	Double Dose
>5 - 10	0.5 tablet	1 tablet
>10 -20	1 tablet	2 tablets

The dose may be doubled, still administered once daily, to a minimum dose of 0.5 mg/kg (range 0.5-1.0), if judged clinically necessary and advised by the veterinary surgeon.

Cats:

FORTEKOR Flavour should be administered orally at a minimum dose of 0.5 mg (range 0.5-1.0) benazepril hydrochloride/kg body weight once daily according to the following table:

Weight of Cat (kg)	Fortekor® 2.5 mg
2.5 - 5	0.5 tablet
>5 - 10	1 tablet

Overdose (symptoms, emergency procedures, antidotes), if necessary

FORTEKOR Flavour reduced erythrocyte counts in normal cats when dosed at 10 mg/kg body weight once daily for 12 months and in normal dogs when dosed at 150 mg/kg body weight once daily for 12 months, but this effect was not observed at the recommended

dose during clinical trials in cats or dogs.

Transient reversible hypotension may occur in cases of accidental overdose. Therapy should consist of intravenous infusion of warm isotonic saline.

Withdrawal period

Not applicable.

Pharmacological particulars

Pharmacotherapeutic group: ACE Inhibitors, plain.

ATC vet code: QC09AA07.

Pharmacodynamic properties

Benazepril hydrochloride is a prodrug hydrolysed *in vivo* to its active metabolite, benazeprilat. Benazeprilat is a highly potent and selective inhibitor of ACE, thus preventing the conversion of inactive angiotensin I to active angiotensin II and thereby also reducing synthesis of aldosterone. Therefore, it blocks effects mediated by angiotensin II and aldosterone, including vasoconstriction of both arteries and veins, retention of sodium and water by the kidney and remodelling effects (including pathological cardiac hypertrophy and degenerative renal changes).

FORTEKOR Flavour causes long-lasting inhibition of plasma ACE activity in dogs and cats, with more than 95% inhibition at peak effect and significant activity (>80% in dogs and >90% in cats) persisting 24 hours after dosing.

FORTEKOR Flavour reduces the blood pressure and volume load on the heart in dogs with congestive heart failure.

In cats with experimental renal insufficiency, FORTEKOR Flavour normalized the elevated glomerular capillary pressure and reduced the systemic blood pressure.

Reduction in glomerular hypertension may retard the progression of kidney disease by inhibition of further damage to the kidneys. Placebo controlled clinical field studies in cats with chronic kidney disease (CKD) have demonstrated that FORTEKOR Flavour significantly reduced levels of urine protein and urine protein to creatinine ratio (UPC); this effect is probably mediated via reduced glomerular hypertension and beneficial effects on the glomerular basement membrane.

No effect of FORTEKOR Flavour on survival in cats with CKD has been shown, but FORTEKOR Flavour increased the appetite of the cats, particularly in more advanced cases.

Pharmacokinetic particulars

After oral administration of benazepril hydrochloride, peak levels of benazepril are attained rapidly (T_{max} 0.5 hour in dogs and within 2 hours in cats) and decline quickly as the active substance is partially metabolised by liver enzymes to benazeprilat. The systemic bioavailability is incomplete (~13% in dogs) due to incomplete absorption (38% in dogs, <30% in cats) and first pass metabolism.

In dogs, peak benazeprilat concentrations (C_{max} of 37.6 ng/ml after a dose of 0.5 mg/kg benazepril hydrochloride) are achieved with a T_{max} of 1.25 hours.

In cats, peak benazeprilat concentrations (C_{max} of 77.0 ng/ml after a dose of 0.5 mg/kg

benazepril hydrochloride) are achieved with a T_{max} of 2 hours.

Benazeprilat concentrations decline biphasically: the initial fast phase (t_{1/2}=1.7 hours in dogs and t_{1/2}=2.4 hours in cats) represents elimination of free drug, while the terminal phase (t_{1/2}=19 hours in dogs and t_{1/2}=29 hours in cats) reflects the release of benazeprilat that was bound to ACE, mainly in the tissues. Benazepril and benazeprilat are extensively bound to plasma proteins (85-90%), and in tissues are found mainly in the liver and kidney.

There is no significant difference in the pharmacokinetics of benazeprilat when benazepril hydrochloride is administered to fed or fasted dogs. Repeated administration of FORTEKOR Flavour leads to slight bioaccumulation of benazeprilat (R=1.47 in dogs and R=1.36 in cats with 0.5 mg/kg), steady state being achieved within a few days (4 days in dogs).

Benazeprilat is excreted 54% via the biliary and 46% via the urinary route in dogs and 85% via the biliary and 15% via the urinary route in cats. The clearance of benazeprilat is not affected in dogs or cats with impaired renal function and therefore no adjustment of FORTEKOR Flavour dose is required in either species in cases of renal insufficiency

Pharmaceutical particulars

List of excipients

Cellulose microcrystalline
Crospovidone
Povidone
Basic butylated methacrylate copolymer
Silicon dioxide anhydrous
Sodium laurilsulphate
Dibutyl sebacate
Silica colloidal anhydrous
Stearic acid
Yeast powder
Artificial powdered beef flavour

Incompatibilities

Not applicable.

Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years

Shelf-life of tablet halves: 2 days

Special precautions for storage

Do not store above 25 °C.

Each time an unused half tablet is stored, it should be returned to the open blister space, inserted back into the cardboard box and kept in a safe place out of the sight and reach of children.

Nature and composition of immediate packaging

14 tablets per aluminium/aluminium blister. Cardboard box with:

1 blister (14 tablets)

2 blisters (28 tablets)

4 blisters (56 tablets)

10 blisters (140 tablets)

Not all pack sizes may be marketed.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

Marketing Authorisation Holder (if different from distributor)

NA

Marketing Authorisation Number

Vm 00879/4045

Significant changes

Date of the first authorisation or date of renewal

28 April 2010

Date of revision of the text

March 2016

Any other information

Legal category

Legal category:POM-V

GTIN

GTIN description:FORTEKOR FLAVOUR 5MG TABLETS FOR CATS AND DOGS 28 pack

GTIN:05037694022259

GTIN description:FORTEKOR FLAVOUR 5MG TABLETS FOR CATS AND DOGS 56 pack

GTIN:05037694023737

Elanco UK AH Limited

Telephone:01256 353131 (24 hr response) Veterinary Technical Services

Website:www.myelanco.co.uk

Email:elancouk@elanco.com