

FACT SHEET

Laminitis - J.T Ferrie F.W.C.F.

PAGE 1



equinevetclinic.co.uk

Laminitis is a disease associated with ischaemia of digital epidermal tissues. The bond between the dermal and epidermal laminae (the inter-laminar bond) is the only means of support of the coffin bone within the hoof. Horses are suspended within their hooves by this interlaminar bond and if the laminar bond is destroyed; the pedal bone moves distally within the hoof capsule.

Laminitis can be Acute (actively Lamanitic) or Chronic (had laminitis in the past but not painful presently).

Laminitis can be: **Rotational** (the laminar bond has broken down at the dorsal region of the hoof and the coffin bone is pulled downwards by the deep flexor tendon); or **Sinking** (where a greater proportion of the Laminar bed is deprived of blood, and consequently nutrients, resulting in the coffin bone sinking downwards within the hoof capsule).

Common Causes

Obesity

The most common type seen in the UK. Many horses and ponies do little work. Unless careful calorie controlled diets are provided these animals very easily become overweight. Native ponies and cobs are very efficient at metabolising quite low energy forage and are unable to cope with fertilized cattle pasture.

Toxaemia

Toxic conditions such as Metritis (retained afterbirth) put the horse at high risk of developing laminitis.

Mechanical

Fast or prolonged work on hard surfaces may induce laminitis also following a non-weight bearing lameness the contra lateral limb may founder.

Hormonal

Some laminitis cases appear to be hypothyroid related.

Cushings Disease

Symptoms of Cushings disease in elderly animals include an enlarged pituitary gland or a pituitary tumour which can cause the adrenal gland to release abnormal amounts of cortisol into the bloodstream and increased levels of cortisol often leads to the animal developing Laminitis.

Diabetes

Variations in insulin levels often seen in diabetes (metabolic syndrome) are thought to induce laminitis.

Stress


Any stress such as overworking unfit horses, prolonged travelling in hot (or cold) conditions, may result in laminitis in some animals. Heat in the feet; is a very unreliable diagnostic indicator. Foot temperature normally varies throughout the day.

Bio Mechanical Forces

The downward force of the body weight is neutralised by the resisting force of the ground. The primary forces involved are the pulling force of the deep digital flexor tendon, and the tearing force at the junction of the white line and the sole




an actively laminitic horse




x-ray showing seroma and sinking



morbid specimen showing total irreparable laminar damage



Venogram showing no blood supply to the Dorsal Laminae



Bio Mechanical Forces

FACT SHEET

Laminitis - J.T Ferrie F.W.C.F.

PAGE 2

Treatment

The condition is always life threatening, there is no such thing as a 'touch' of laminitis. Therefore Veterinary advice should be sought immediately. Usually, non steroidal anti-inflammatory drugs will be prescribed, in conjunction with frog support, along with the removal of the cause (if known).

Farriery treatment normally follows the abatement of the symptoms and fresh x-rays are advisable to locate the true position of the bone inside the hoof capsule. A heart bar is my shoe of choice, usually made of steel or an alloy and nailed on lightly. This shoe offers stability to the bone column of the leg and hoof.

New technology allows us to glue hydroplastic heart bars to feet, should nailing prove too traumatic.

Hereditary predisposition to laminitis in this country is unproven. However families of animals often have the same owner whose predisposition to recurring poor management is certainly proven.

Laminitis does not just affect the front feet. Although due to 65% of bodyweight being carried on the forelimbs, the front feet are usually more badly affected.

Fitting a Heartbar Shoe to a Lamanitic Horse

The correct method of application of a Heartbar is crucial to the recovery of a Lamanitic animal. This should always be done with Veterinary consultation.

Current x-rays are essential in positioning the shoe correctly.

The shoe is designed so that the frog carries a significant proportion of weight that would have been normally supported by the wall and sole; especially at the heels. The Farriery term for this is a loaded Heartbar.

Loaded Heartbar.

The bar in the shoe replicates the triangular shape of the frog, and lays flat on it within its perimeter. The tip of the bar must be perpendicular to a line drawn from the Extensor Process of the Coffin bone within the hoof capsule. The drawing pin shows the true point of the frog, giving an external reference point..

If the bar in the shoe is positioned too far forward the arterial blood supply to the Laminae and sole will be compromised usually resulting in an abscess. If too far back the tip of the bone is driven downwards and can protrude through the sole.

The x-rays will not only show the current position of the Coffin bone in relation to the hoof capsule, but can also show any pockets of serum which have filled the void created by the dead Laminae.

When fitting a loaded Heartbar any serum or gas pockets should be opened to release the Hydrostatic pressure within the hoof and offer relief to the horse. It should be remembered that this breakdown of the Laminar bond occurs in three dimensions. Often the bone will displace medially due to the centre of weight bearing being carried down the inside of the limb.


Shoeing the Chronic Lamanitic Horse

When turning in a tight circle is not problematic the heartbars may be removed and replaced with shoes which don't impede the distended laminae at the toe ie.either unclipped or side clipped shoes.

The prognosis of a full recovery is always guarded, however, if the source can be found and managed, then with proper regular Farriery and Veterinary management it is not unreasonable to expect a return to limited work.



concave heartbar



the tip of the coffin bone coming through the sole.



medial displacement of the hoof capsule

